arXiv:1410.5555v1 [cs.CC] 21 Oct 2014

On computational complexity of length embeddability of graphs

Mikhail Tikhomirov

Abstract

A graph G is embeddable in $\mathbb{R}^d$ if vertices of G can be assigned with points of $\mathbb{R}^d$ in such a way that all pairs of adjacent vertices are at the distance 1. We show that verifying embeddability of a given graph in $\mathbb{R}^d$ is NP-hard in the case $d > 2$ for all reasonable notions of embeddability.

1 Introduction

The *distance graph of* $S \subset \mathbb{R}^d$ is defined as the graph $G = (V, E)$, where $V = S$ and E is the set of all pairs of points $x, y \in S$ such that x and y are at the distance 1. A graph is a *distance graph in* $\mathbb{R}^d$ if it is isomorphic to the distance graph of some set $S \subset \mathbb{R}^d$. Some famous problems concerning distance graphs are the Erdős' unit distance problem on the maximal number of unit distances between n points in $\mathbb{R}^2$ (see [1], [2], [3]), the Hadwiger–Nelson problem on the chromatic number of $\mathbb{R}^2$ (see [1], [4], [5]), etc.; surveys of various results about distance graphs can be found at [6], [7].

We also consider a similar notion of *embeddability in* $\mathbb{R}^d$ (see, e.g., [8]). A graph $G = (V, E)$ is *embeddable in* $\mathbb{R}^d$ if there exists a mapping $\varphi : V \to \mathbb{R}^d$ such that $||\varphi(u) - \varphi(v)||_{\mathbb{R}^d} = 1$ for all pairs $(u, v) \in E$. It is clear that any distance graph in $\mathbb{R}^d$ is embeddable in $\mathbb{R}^d$ but the converse does not always hold. These two notions differ in the following:

- Different vertices of an embeddable graph may be assigned with the same point in $\mathbb{R}^d$ while all vertices of a distance graph should be assigned with pairwise distinct points.
- Non-adjacent vertices of an embeddable graph can be located at the distance 1 while non-adjacent vertices of a distance graph are forbidden to be placed at distance 1.

We will say that an embedding $\varphi : V \to \mathbb{R}^d$ is *strict* if $\forall u, v \in V\ (u, v) \in E \iff ||\varphi(u) - \varphi(v)||_{\mathbb{R}^d} = 1$; we will say that an embedding $\varphi : V \to \mathbb{R}^d$ is *injective* if $\forall u, v \in V\ v \neq u \Rightarrow \varphi(v) \neq \varphi(u)$. It is clear that a graph G is a distance graph in $\mathbb{R}^d$ iff there exists a strict and injective embedding of G in $\mathbb{R}^d$. Thus we obtain four different notions of embeddability (strict/non-strict, injective/non-injective) which include two notions described above.

For each of the four notions of embeddability in $\mathbb{R}^d$ we can pose the computational decision problem of determining embeddability of chosen type for the given graph; we shall call this problem **$\mathbb{R}^d$-UNIT-DISTANCE-(STRICT)-(INJECTIVE)-EMBEDDABILITY** depending on the embeddability type. The computational complexity of these problems is studied in [8], [9]. In [9] it is shown that **$\mathbb{R}^d$-UNIT-DISTANCE-(STRICT)-(INJECTIVE)-EMBEDDABILITY** is NP-hard for each type of embeddability and each value of $d \geqslant 2$. Unfortunately, the proof in [9] for the case $d > 2$ is false as it is based on the result [10] due to Lovász which states the upper bound $d+1$ for the chromatic number of the d-dimensional sphere circumscribed about a regular simplex on $d+1$ vertices with unit length edges. In [11], [12] Raigorodskii points out that this bound is wrong and proves an exponential lower bound of this value; thus a new proof is needed for the case $d > 2$, which is the point of this paper.

The main result is

Theorem 1. *Computational problems $\mathbb{R}^d$-UNIT-DISTANCE-EMBEDDABILITY, $\mathbb{R}^d$-UNIT-DISTANCE-STRICT-EMBEDDABILITY, $\mathbb{R}^d$-UNIT-DISTANCE-INJECTIVE-EMBEDDABILITY, $\mathbb{R}^d$-UNIT-DISTANCE-STRICT-INJECTIVE-EMBEDDABILITY are NP-hard for each $d > 2$.*

To prove this result we contrust a reduction of the classic NP-complete problem of graph vertex 3-coloring (**3-COLORING**) (see [13]) to each of the four embeddability problems: for any given graph G we explicitly construct a graph H = 3-COLORING-$\mathbb{R}^d$-UNIT-DISTANCE-EMBEDDABILITY-REDUCTION(G) such that the size of H is linear in the size of G (for every fixed d) and the following conditions hold:

- If no valid vertex 3-coloring of G exists, then there is no embedding of H in $\mathbb{R}^d$;
- If a valid vertex 3-coloring of G exists, then there is a strict injective embedding of H in $\mathbb{R}^d$.

The possibility of such construction implies NP-hardness of all four mentioned problems. It should be mentioned that the question whether the described problems lie in NP is open.

2 Notion of rod

Let us introduce some necessary definitions.

A *weighted graph* $G = (V, E, w)$ is an ordered triple such that (V, E) is a graph and $w : E \to \mathbb{R}_+$ is a function that assigns a positive number to each element of E; for every edge $e \in E$ we will say that $w(e)$ is the *length* of the edge e. If $w \equiv 1$, the weighted graph G is called a *unit distance graph*. A *length embedding* (or, more simply, an *embedding*) of the weighted graph $G = (V, E, w)$ in $\mathbb{R}^d$ is a map $\varphi : V \to \mathbb{R}^d$ such that $\forall u, v \in V\ (u, v) \in E \Rightarrow ||\varphi(u) - \varphi(v)||_{\mathbb{R}^d} = w((u, v))$.

***Remark**: In the sequel, we will identify vertices of the graph with points of $\mathbb{R}^d$ — their images under the embedding if that doesn't cause confusion.*

An embedding φ of the weighted graph $G = (V, E, w)$ in $\mathbb{R}^d$ is called *non-critical* if the following conditions hold:

- $\forall u, v \in V\ u \neq v \Rightarrow \varphi(u) \neq \varphi(v)$ (no two vertices are at the same point);
- $\forall u, v \in V\ (u, v) \notin E \Rightarrow ||\varphi(u) - \varphi(v)||_{\mathbb{R}^d} \neq 1$ (no two non-adjacent vertices are at the distance 1);
- no three vertices lie on a (one-dimensional) straight line.

Consider a weighted graph $G = (V, E, w)$ and a pair of its vertices $u, v \in V$. The graph G is called a *(d-dimensional) (u, v)-rod of length l* if the following conditions hold:

- the distance between vertices u and v is equal to l in each embedding of G in $\mathbb{R}^d$;
- there exists a non-critical embedding of G in $\mathbb{R}^d$.

If a unit distance graph G is also a d-dimensional (u, v)-rod of length l, we call G a *d-dimensional unit distance (u, v)-rod of length l*.

A weighted graph $G = (V, E, w)$ is called a *(unit distance) d-dimensional rod of length l* if there exist two vertices $u, v \in V$ such that G is a (unit distance) d-dimensional (u, v)-rod of length l.

We suppose that d is a fixed constant throughout the whole paper, thus in the sequel we will write "rod" for "d-dimensional rod".

Lemma 1. *Let $G = (V_G, E_G, w_G)$, $H = (V_H, E_H, w_H)$ be weighted graphs. Suppose $V_G \cap V_H = \{u, v\}$, $e = (u, v) \in E(G)$, $w_G(e) = l$, and H is a (u, v)-rod of length l. Let $G' = (V_{G'}, E_{G'}, w_{G'})$, where $V_{G'} = V_G \cup V_H$, $E_{G'} = (E_G \backslash \{e\}) \cup E_H$, $w_{G'} = w_G I(E_G \backslash \{e\}) + w_H I(E_H)$ (informally, we replace the edge e in G by the subgraph H to obtain G'). Then:*

- *If there is no embedding of G in $\mathbb{R}^d$, then there is no embedding of G' in $\mathbb{R}^d$.*
- *If there exists a **non-critical** embedding of G in $\mathbb{R}^d$, then there exists a **non-critical** embedding of G' in $\mathbb{R}^d$.*

Proof. In any embedding of G' the distance between vertices u and v is equal to l. Suppose we have an embedding of G'; we can erase all vertices outside V_G to obtain an embedding of G. The first claim is thus proven.

Now consider a non-critical embedding φ_G of the weighted graph G in $\mathbb{R}^d$. Construct an embedding $\varphi_{G'}$ of G' as follows:

- Let $\varphi_{G'}(x) = \varphi_G(x)$ for all $x \in V_G$;
- Choose a non-critical embedding φ_H of the weighted graph H such that $\varphi_H(u) = \varphi_G(u)$, $\varphi_H(v) = \varphi_G(v)$ (such embedding exists since $||\varphi_G(u) - \varphi_G(v)|| = l$ and H is a (u, v)-rod of length l); let $\varphi_{G'}(y) = \varphi_H(y)$ for all $y \in V_H$.

It is clear that this definition of $\varphi_{G'}$ is consistent. However, it is possible that $\varphi_{G'}$ is not a non-critical embedding. Note that no vertex of $V_{G'}\backslash\{u,v\}$ lies on the straight line uv since the embeddings φ_G and φ_H are non-critical.

Let S denote the set of all rotations of $\mathbb{R}^d$ about the line uv. S is isomorphic to the $(d-2)$-dimensional sphere (each rotation can be assigned with the image of some point which doesn't lie on uv). For any $\psi \in S$ let $\psi *_H \varphi_{G'}$ denote the mapping from $V_{G'}$ in $\mathbb{R}^d$ such that $\psi *_H \varphi_{G'}(x) = \varphi_{G'}(x)$ for every $x \in V_G$ and $\psi *_H \varphi_{G'}(y) = \psi(\varphi_{G'}(y))$ for every $y \in V_H$; clearly, this definition is consistent. It is also clear that for every rotation $\psi \in S$ the mapping $\psi *_H \varphi_{G'}$ is an embedding of G' in $\mathbb{R}^d$.

We now show that there exists a rotation $\psi \in S$ such that $\psi *_H \varphi_{G'}$ is a non-critical embedding of G' in $\mathbb{R}^d$. Consider all $\psi \in S$ such that the embedding $\psi *_H \varphi_{G'}$ is not non-critical for some reason. In that case, one of the following conditions must hold:

- The embedding $\psi *_H \varphi_{G'}$ places two vertices of G' (denote them x and y) at the same point. It follows from the non-criticality of φ_G and φ_H that x and y cannot lie both in V_G or both in V_H. Thus WLOG $x \in V_G\backslash\{u,v\}$, $y \in V_H\backslash\{u,v\}$.

  The vertex y does not lie on the line uv and no two rotations place y at the same point. Therefore for every pair of vertices x, y there is at most one rotation $\psi \in S$ that superposes x and $\psi(y)$, thus the set of all rotations ψ such that the embedding $\psi *_H \varphi_{G'}$ places some two vertices in the same point is finite and its spherical measure in S is zero.

- The embedding $\psi *_H \varphi_{G'}$ places two non-adjacent vertices of G' (denote them x and y once more) at the distance 1. Once again, $x, y \in V_G$ or $x, y \in V_H$ leads to a contradiction; thus WLOG $x \in V_G\backslash\{u,v\}$, $y \in V_H\backslash\{u,v\}$.

  Let P_y denote the $(d-2)$-dimensional sphere — the locus of the point $\psi(y)$ for all $\psi \in S$; the radius of P_y is non-zero since y does not lie on the line uv. If $||x - \psi(y)||_{\mathbb{R}^d} = 1$, then $\psi(y)$ lies on the $(d-1)$-dimensional sphere of radius 1 centered at x; denote it P_x. We assume that the intersection of P_x and P_y is not empty.

  If P_x contains P_y as a subset, then x must lie on the line uv; that would contradict the non-criticality of φ_G. Otherwise, the intersection of P_x and P_y is a $(d-3)$-dimensional sphere (possibly, of zero radius).

  In any case, the set of rotations that place x and $\psi(y)$ at the distance 1 has zero measure in S. Thus the set of rotations that place some two non-adjacent vertices at the distance 1 has zero measure in S.

- The embedding $\psi *_H \varphi_{G'}$ places some three vertices on a straight line; denote these vertices x, y, z. Similarly to previous cases, if we assume $x, y, z \in V_G$ or V_H we arrive at a contradiction.

  WLOG, let $x, y \in V_G\backslash\{u,v\}$, $z \in V_H\backslash\{u,v\}$. Since the point z can not lie on the line uv, the sphere $\psi(z)$ for $\psi \in S$ has non-zero radius and the line xy passes through the point $\psi(z)$ for at most two values of ψ.

  Now let $x \in V_G\backslash\{u,v\}$ and $y, z \in V_H\backslash\{u,v\}$. The rotation $\psi \in S$ places the point x on the line $\psi(yz)$ iff the point $\psi^{-1}(x)$ lies on the line yz (here ψ^{-1} means

the inverse rotation of ψ), therefore in this case the line yz must cross the locus of $\psi^{-1}(x)$ for all $\psi \in S$. Clearly, the locus is a sphere of non-zero radius, thus line $\psi(yz)$ passes through the point x for at most two values of ψ.

It follows from the above that the set of rotations $\psi \in S$ such that $\psi *_H \varphi_{G'}$ places some three vertices on a straight line is finite.

To sum up, the set of rotations ψ such that the embedding $\psi *_H \varphi_{G'}$ is not non-critical has zero measure in the $(d-2)$-dimensional sphere of all possible rotations about the line uv. Therefore almost every rotation $\psi \in S$ yields a non-critical embedding $\psi *_H \varphi_{G'}$ of the graph G' in $\mathbb{R}^d$.

□

3 Construction of rods

Let $h = \sqrt{\frac{d+1}{2d}}$ denote the altitude length of a regular d-dimensional simplex with the edge length 1; denote $D = 2h$. Clearly, $D > \sqrt{2}$.

Lemma 2. *Let G and H be unit distance rods of length a and b respectively. Then there exists a unit distance rod of length ab.*

Proof. It suffices to make lengths of all edges of G be equal to b and successively apply Lemma 1 to every edge of the resulting graph and the graph H.

□

Consider a graph M_d on a set of vertices $V_{M_d} = K_1 \cup K_2 \cup \{A, B, C\}$, $|K_1| = |K_2| = d$. Add the following edges of unit length to M_d:

- make cliques on K_1 and K_2;
- connect the vertices A and B with every vertex of K_1;
- connect the vertices A and C with every vertex of K_2;
- finally, connect the vertices B and C.

The graph M_d is called a *d-dimensional Moser spindle* (the figure 1 illustrates a 5-dimensional Moser spindle). Is it easy to see that M_d is a unit distance d-dimensional (A, B)-rod of length D.

Repeatedly applying Lemma 2 to copies of M_d, we arrive at

Corollary 1. *For every non-negative integer k there exists a unit distance d-dimensional rod of length D^k.*

Lemma 3. *For all numbers a, b such that $0 < a < b < 1$ there exists a number l satisfying $a < l < b$ and a graph G such that G is a unit-distance d-dimensional rod of length l.*



Figure 1: 5-dimensional Moser spindle

Proof. Construct G as follows. Choose a set of vertices K of size $d-1$ and connect its elements pairwise by unit length edges. Then, take a sequence of vertices $v_1, \ldots, v_n$ (the exact number of vertices n will be determined later) and connect every vertex of the sequence v_i with every vertex of K by a unit length edge. If the location of vertices of K is fixed, then all vertices $v_1, \ldots, v_n$ must lie on some circle centered at O, where O is the center of the regular simplex with vertices in K. The radius of the circle is equal to the altitude length of the $(d-1)$-face of the regular d-simplex with the side length 1, i.e. $\sqrt{\frac{d}{2(d-1)}} = r$. Let π denote the plane containing this circle.

For each i from 1 to $n-1$ connect the vertices v_i and v_{i+1} by an edge of length 1; also for each i from 1 to $n-2$ connect the vertices v_i and v_{i+2} by an edge of length D. Now in every embedding of the graph G the angle $\angle v_i O v_{i+1}$ is equal to the dihedral angle of a regular d-simplex; denote this angle $\alpha = \arccos \frac{1}{d}$. Additionally, the least rotation of the plane π about the point O that moves the point v_i to v_{i+1} has the same direction for every i. It is clear that no three vertices of G lie on a straight line.

Let us introduce an angular coordinate system ψ on π centered at O such that $\psi(v_1) = 0$, $\psi(v_2) = \alpha$. Clearly, $\psi(v_i) = (i-1)\alpha \mod 2\pi$ (by $\alpha \mod 2\pi$ we mean $\alpha + k \times 2\pi$ for an integer k such that $0 \leqslant \alpha + k \times 2\pi < 2\pi$). By Niven's theorem (see [14], Corollary 3.12), $\alpha/2\pi$ can not be a rational number when $d \geqslant 3$, therefore the infinite sequence $x_i = (i-1)\alpha \mod 2\pi$ is dense in $[0; 2\pi]$. Thus there exists a positive integer N such that $x_N \in (2\arcsin\frac{a}{2r}; 2\arcsin\frac{b}{2r})$ and $||v_1 - v_N|| \in (a; b)$.

It follows from the above that the graph G is a d-dimensional (v_1, v_N)-rod. Finally, successively apply Lemma 2 to each D-length edge of the graph G and the graph M_d; the resulting graph is a unit distance d-dimensional (v_1, v_N)-rod that satisfies all the conditions.

□

Theorem 2. *For all numbers a, b such that $0 < a < b$ there exists a number l satisfying $a < l < b$ and a graph G such that G is a unit-distance d-dimensional rod of length l.*

Proof. Choose a non-negative integer k such that $D^k > b$ and denote G' the rod obtained by applying Lemma 3 for numbers $\frac{a}{D^k}$ and $\frac{b}{D^k}$. Now apply Lemma 2 to the graph G' and the rod of length D^k.

□

Let RodLength(a, b) denote the number l produced by Theorem 2 for given numbers a and b, and Rod(a, b) denote the rod of corresponding length.

4 The reduction setup

Consider a graph $G = (V_G, E_G)$ — the input of the **3-COLORING** problem. We now construct a weighted graph $H = (V_H, E_H, w_H) = $ 3-COLORING-$\mathbb{R}^d$-EMBEDDABILITY-REDUCTION(G) such that the embeddability of H in $\mathbb{R}^d$ is equivalent to the existence of a solution to the **3-COLORING** for the graph G. We shall idenitify the elements of V_G and the integers from 1 to $|V_G|$ for the sake of convenience.

To establish properties of the following setup we will need the following

Lemma 4. *Let $0 \leqslant l < L \leqslant R < r$, $\delta = \min(L - l, r - R)$.*
Let also $G = (V_G, E_G, w_G)$, $H = (V_H, E_H, w_H)$ — weighted graphs,
$v, u \in V_G$, $V_H = V_G \sqcup \{z\}$, $E_H = E_G \sqcup \{(v, z), (u, z)\}$,
$w_H(e) = w_G(e)$ for all $e \in E_G$,
$w_H((v, z)) = a \in \left(\frac{L+R}{2} - \frac{\delta}{3}; \frac{L+R}{2} + \frac{\delta}{3}\right)$,
$w_H((u, z)) = b \in \left(\frac{R-L}{2} + \frac{\delta}{3}; \frac{R-L}{2} + \frac{\delta}{2}\right)$.
Then:

- *In every embedding of the graph H the inequalities $l < ||v - u|| < r$ hold.*
- *If there exists a* non-critical *embedding of G such that $L \leqslant ||v - u|| \leqslant R$, then there exists a* non-critical *embedding of H.*

Proof. First of all, let us show that $l < a - b < L \leqslant R < a + b < r$. Indeed,

$$a - b > \left(\frac{L+R}{2} - \frac{\delta}{3}\right) - \left(\frac{R-L}{2} + \frac{\delta}{2}\right) > L - \delta \geqslant l;$$

$$a - b < \left(\frac{L+R}{2} + \frac{\delta}{3}\right) - \left(\frac{R-L}{2} + \frac{\delta}{3}\right) = L;$$

$$a + b > \left(\frac{L+R}{2} - \frac{\delta}{3}\right) - \left(\frac{R-L}{2} + \frac{\delta}{3}\right) = R;$$

$$a + b < \left(\frac{L+R}{2} + \frac{\delta}{3}\right) + \left(\frac{R-L}{2} + \frac{\delta}{2}\right) < R + \delta \leqslant r.$$

Consider any embedding of the graph H. It follows from the triangle inequality applied to vertices v, z, u that $l < a - b \leqslant ||u - v|| \leqslant a + b < r$. The first claim is thus proven.

Now, consider a non-critical embedding of the graph G such that $||u - v|| \in [L; R]$. It follows from $a - b < L \leqslant ||u - v|| \leqslant R < a + b$ that it is possible to place the vertex z in such a way that $||v - z|| = a$, $||u - z|| = b$ and z does not lie on the line vu. We have obtained an embedding of the graph H; it is possible to modify this embedding to obtain a non-critical embedding by choosing an appropriate rotation of z about the

line vu; the proof of the existence of such rotation copies the similar proof from Lemma 1 almost entirely.

□

Denote $r_0 = \sqrt{\frac{d}{2(d-1)}}$, $\text{chord}(\alpha) = 2r_0 \sin \alpha/2$ — the length of the chord which contracts an α-measured arc of a circle of radius r_0, $\varepsilon = \frac{\pi}{24}$.

Let us introduce additional notation as follows:

$\delta_{uu} = \min(\text{chord}(2\pi/3) - \text{chord}(2\pi/3 - \varepsilon/2), \text{chord}(2\pi/3 + \varepsilon/2) - \text{chord}(2\pi/3))$,
$a_{uu} = \text{RodLength}(\text{chord}(2\pi/3) - \delta_{uu}/3, \text{chord}(2\pi/3) + \delta_{uu}/3)$,
$b_{uu} = \text{RodLength}(\delta_{uu}/3, \delta_{uu}/2)$,
$\delta_{uv} = \text{chord}(\pi/3 - \varepsilon/2) - \text{chord}(\pi/3 - \varepsilon)$,
$a_{uv} = \text{RodLength}\left(\frac{2r_0 + \text{chord}(\pi/3-\varepsilon/2)}{2} - \delta_{uv}/3, \frac{2r_0 + \text{chord}(\pi/3-\varepsilon/2)}{2} + \delta_{uv}/3\right)$,
$b_{uv} = \text{RodLength}\left(\frac{2r_0 - \text{chord}(\pi/3-\varepsilon/2)}{2} + \delta_{uv}/3, \frac{2r_0 - \text{chord}(\pi/3-\varepsilon/2)}{2} + \delta_{uv}/2\right)$,
$\delta_{vv} = \text{chord}\left(\frac{2}{3}\pi - \varepsilon\right) - \text{chord}\left(\frac{5}{2}\varepsilon\right)$,
$a_{vv} = \text{RodLength}\left(\frac{2r_0 + \text{chord}(2\pi/3-\varepsilon)}{2} - \delta_{vv}/3, \frac{2r_0 + \text{chord}(2\pi/3-\varepsilon)}{2} + \delta_{vv}/3\right)$,
$b_{vv} = \text{RodLength}\left(\frac{2r_0 - \text{chord}(2\pi/3-\varepsilon)}{2} + \delta_{vv}/3, \frac{2r_0 - \text{chord}(2\pi/3-\varepsilon)}{2} + \delta_{vv}/2\right)$.

Construct H = 3-COLORING-$\mathbb{R}^d$-EMBEDDABILITY-REDUCTION(G) as follows:

$$V_H = K \cup U \cup V \cup Aux, E_H = E_K \cup E_{KU} \cup E_{KV} \cup E_U \cup E_{VU} \cup E_V.$$

Here:

- Aux is the set of all auxiliary vertices used in the sequel of the description ($aux_{\dots}$);
- K — the set of vertices of size $d-1$;
  E_K — the set of edges connecting all pairs of vertices of K;
- $U = \{u_0, u_1, u_2\}$;
  E_{KU} — the set of edges connecting every vertex of U with every vertex of K;
  $E_U = \{(u_0, aux_{u_0,u_1}), (aux_{u_0,u_1}, u_1), (u_0, aux_{u_0,u_2}), (aux_{u_0,u_2}, u_2), (u_1, aux_{u_1,u_2}), (aux_{u_1,u_2}, u_2)\}$;
- $V = \{v_1, \dots, v_{|V_G|}\}$;
  E_{KV} — the set of edges connecting every vertex of V with every vertex of K;
  $E_{VU} = \bigcup\limits_{v \in V} \bigcup\limits_{u \in U} \{(v, aux_{v,u}), (aux_{v,u}, u)\}$;
  $E_V = \bigcup\limits_{v_i, v_j : (i,j) \in E_G} \{(v_i, aux_{v_i,v_j}), (aux_{v_i,v_j}, v_j)\}$.

The edge lengths are assigned as follows:

- $e \in E_K \cup E_{KU} \cup E_{KV} \Rightarrow w_H(e) = 1$;
- $w_H(u_0, aux_{u_0,u_1}) = w_H(u_0, aux_{u_0,u_2}) = w_H(u_1, aux_{u_1,u_2}) = a_{uu}$,
  $w_H(aux_{u_0,u_1}, u_1) = w_H(aux_{u_0,u_2}, u_2) = w_H(aux_{u_1,u_2}, u_2) = b_{uu}$;

- for all $v \in V$, $u \in U$: $w_H(v, aux_{v,u}) = a_{uv}$, $w_H(aux_{v,u}, u) = b_{uv}$;
- for all pairs $v_i, v_j \in V$ such that $(i,j) \in E_G$: $w_H(v_i, aux_{v_i,v_j}) = a_{vv}$, $w_H(aux_{v_i,v_j}, v_j) = b_{vv}$.

Theorem 3. *Let $G = (V_G, E_G)$ and H = 3-COLORING-$\mathbb{R}^d$-EMBEDDABILITY-REDUCTION(G) described above. Then:*

- *If there is no valid 3-coloring of vertices of G, then there is no embedding of H in $\mathbb{R}^d$.*
- *If a valid 3-coloring of vertices of G exists, then there exists a non-critical embedding of H in $\mathbb{R}^d$.*

Proof. Consider an embedding of the graph H in $\mathbb{R}^d$; let us construct a valid 3-coloring of vertices of G. In every embedding of H all vertices of the set $V \cup U$ are located on some circle of radius r_0; denote this circle ρ and its center O.

Successively apply the first part of Lemma 4 with the following parameters.

- Let $v = u_0$, $u = u_1$, $z = aux_{u_0,u_1}$, $l = \text{chord}(2\pi/3 - \varepsilon/2)$, $L = R = \text{chord}(2\pi/3)$, $r = \text{chord}(2\pi/3 + \varepsilon/2)$.
  We obtain that $\text{chord}(2\pi/3 - \varepsilon/2) < ||u_0 - u_1|| < \text{chord}(2\pi/3 + \varepsilon/2)$, which is equivalent to $2\pi/3 - \varepsilon/2 < \angle u_0 O u_1 < 2\pi/3 + \varepsilon/2$. We can establish similar inequalities for u_0, u_2 and u_1, u_2.
- Let $v \in V$, $u \in U$, $z = aux_{v,u}$, $l = \text{chord}(\pi/3 - \varepsilon)$, $L = \text{chord}(\pi/3 - \varepsilon/2)$, $R = 2r_0$, $r = \infty$.
  Then $\text{chord}(\pi/3 - \varepsilon) < ||v - u||$, which is equivalent to $\pi/3 - \varepsilon < \angle vOu$.
- Let $v_i, v_j \in V$, $(i,j) \in E_G$: $v = v_i$, $u = v_j$, $z = aux_{v_i,v_j}$, $l = \text{chord}\left(\frac{5}{2}\varepsilon\right)$, $L = \text{chord}(2\pi/3 - \varepsilon)$, $R = 2r_0$, $r = \infty$.
  Then $\text{chord}\left(\frac{5}{2}\varepsilon\right) < ||v_i - v_j||$, which is equivalent to $\frac{5}{2}\varepsilon < \angle v_i O v_j$.

Construct the coloring of vertices of G as follows: if the vertex v_i lies on the shortest arc between u_0 and u_1 in the embedding of H, the vertex $i \in V_G$ is assigned with the color $c(i) = 2$; if v_i lies on the shortest arc between u_0 and u_2, then $c(i) = 1$; otherwise, $c(i) = 0$. It is clear that this coloring is unambigiously defined for any embedding of H. We now prove that this coloring of vertices of G is valid, that is, for every edge $(i,j) \in E(G)$ we have $c(i) \neq c(j)$.

Let us show that for every edge $(i,j) \in E_G$ the shortest arc of ρ between the points v_i and v_j contains at least one vertex of U. Assume the contrary, then WLOG both vertices v_i and v_j lie on the shortest arc between u_0 and u_1, and $\angle u_0 O u_1 = \angle u_0 O v_i + \angle v_i O v_j + \angle v_j O u_1 > (\pi/3 - \varepsilon) + \frac{5}{2}\varepsilon + (\pi/3 - \varepsilon) = 2\pi/3 + \varepsilon/2$. But that contradicts with $\angle u_0 O u_1 < 2\pi/3 + \varepsilon/2$, thus at least one vertex of U must lie between v_i and v_j. In that case the colors of i and j are different; therefore the coloring is valid. The first part of Theorem 3 is thus proven.

Denote $H' = (V_{H'}, E_{H'}, w_{H'})$, where $V_{H'} = K \cup U \cup V$, $E_{H'} = E_K \cup E_{KU} \cup E_{KV}$, $w_{H'} \equiv 1$. Clearly, H' is a subgraph of H.

Now consider a valid vertex 3-coloring of G; let us construct a non-critical embedding of H. First, construct a non-critical embedding of H' as follows:

- choose an arbitrary regular $(d-2)$-simplex with edge length 1 and identify its vertices with vertices of K; let O denote the center of the simplex and ρ denote the locus of all points at the distance 1 from all vertices of the simplex; clearly, ρ is a circle of radius r_0;

- choose an arbitrary equilateral triangle inscribed in ρ; place the vertices u_0, u_1, u_2 at the vertices of the triangle; denote γ_0 the set of all points $x \in \rho$ such that $\angle u_0 O x > \pi - \varepsilon/2$; clearly, γ_0 is an open arc of angular measure ε; similarly define sets γ_1, γ_2;

- suppose the vertex $i \in V_G$ is assigned with color $c(i) \in \{0, 1, 2\}$ in the given 3-coloring; place every vertex $v_i \in V$ in such a way that v_i lies on the arc $\gamma_{c(i)}$ for every $i \in V_G$ and no two vertices of V are at the same point; since the arcs γ_0, γ_1, γ_2 have non-zero angular measure, such arrangement of vertices of V is possible.

It can be easily verified that the arrangement of vertices of $K \cup V \cup U$ described above yields a non-critical embedding of the graph H'.

Now let us add vertices of the set Aux one by one and successively apply the second part of Lemma 4 to show the existence of a non-critical embedding for every new graph. When all vertices of Aux are added, we obtain a non-critical embedding of the graph H since every vertex of Aux is adjacent to exactly two vertices of $V \cup U$.

Successively apply the second part of Lemma 4 with the following parameters.

- Let $v = u_0$, $u = u_1$, $z = aux_{u_0,u_1}$, $l = \mathrm{chord}(2\pi/3 - \varepsilon/2)$, $L = R = \mathrm{chord}(2\pi/3)$, $r = \mathrm{chord}(2\pi/3 + \varepsilon/2)$.
  The points u_0 and u_1 are at the vertices of an equilateral triangle inscribed in the circle ρ, thus $||u_0 - u_1|| = \mathrm{chord}(2\pi/3)$ and the conditions of the lemma are satisfied.
  Apply the lemma in a similar way to u_0, u_2, aux_{u_0,u_2} and u_1, u_2, aux_{u_1,u_2}.

- Let $v \in V$, $u \in U$, $z = aux_{v,u}$, $l = \mathrm{chord}(\pi/3 - \varepsilon)$, $L = \mathrm{chord}(\pi/3 - \varepsilon/2)$, $R = 2r_0$, $r = \infty$.
  There is at least one vertex $u_i \in U$ such that $\angle v O u_i > \pi - \varepsilon/2$, thus $\angle v O u \geqslant |\angle v O u_i - \angle u_i O u| > \pi/3 - \varepsilon/2$ and $||v - u|| > \mathrm{chord}(\pi/3 - \varepsilon/2)$; the conditions of the lemma are satisfied.

- Let $v_i, v_j \in V$, $(i, j) \in E_G$: $v = v_i$, $u = v_j$, $z = aux_{v_i,v_j}$, $l = \mathrm{chord}(\frac{5}{2}\varepsilon)$, $L = \mathrm{chord}(2\pi/3 - \varepsilon)$, $R = 2r_0$, $r = \infty$.
  The points v_i and v_j lie on different arcs $\gamma_{c(i)}$, $\gamma_{c(j)}$. Let u_k denote the vertex of U that lies on the shortest arc between v_i and v_j. Then $\angle v_i O v_j = \angle v_i O u_k + \angle u_k O v_j > 2(\pi/3 - \varepsilon/2) = 2\pi/3 - \varepsilon$, and $||v_i - v_j|| > \mathrm{chord}(2\pi/3 - \varepsilon)$; the conditions of the lemma are satisfied.

After all applications of Lemma 4 we obtain a non-critical embedding of the graph H.

□

The constructed graph H has $O(|V_G| + |E_G|)$ vertices and edges (we recall that the dimension d is a fixed constant), but it contains edges of non-unit length; however, for every such edge its length is equal to $\mathrm{RodLength}(a, b)$ for some a, b; moreover, the set of possible pairs (a, b) is finite and independent on the input graph G. Thus, upon multiple applications of Lemma 2 each edge of non-unit length can be replaced by a subgraph that is isomorphic to $\mathrm{Rod}(a, b)$ for some (a, b); the size of the graph will increase by at most K times, where K is the maximal size of $\mathrm{Rod}(a, b)$ for all used pairs of (a, b); clearly, the value of K depends only on d. Therefore the resulting graph $H' = $ 3-COLORING-$\mathbb{R}^d$-UNIT-DISTANCE-EMBEDDABILITY-REDUCTION(G) has $O(|V_G| + |E_G|)$ vertices and edges as well. Finally, we obtain

Theorem 4. *Let the graph $H' =$ 3-COLORING-$\mathbb{R}^d$-UNIT-DISTANCE-EMBEDDABILITY-REDUCTION(G) be constructed by a given graph $G = (V_G, E_G)$ as described above. Then:*

- *If there is no valid 3-coloring of vertices of G, then there is no embedding of H' in $\mathbb{R}^d$.*
- *If a valid 3-coloring of vertices of G exists, then there exists a non-critical embedding of H' in $\mathbb{R}^d$.*

From Theorem 4, the linearity of the size of H', and the fact that the problem of vertex 3-coloring is NP-hard (see [13]) Theorem 1 eventually follows.

References

[1] P. Brass, W. O. Moser, and J. Pach, *Research problems in discrete geometry*, vol. 18. Springer, 2005.

[2] P. Erdős, "On sets of distances of n points," *American Mathematical Monthly*, pp. 248–250, 1946.

[3] K. B. Chilakamarri, "The unit-distance graph problem: a brief survey and some new results," *Bull. Inst. Combin. Appl*, vol. 8, no. 39, p. C60, 1993.

[4] N. De Bruijn and P. Erdős, "A colour problem for infinite graphs and a problem in the theory of relations," *Indag. Math*, vol. 13, no. 5, pp. 371–373, 1951.

[5] A. M. Raigorodskii, "Borsuk's problem and the chromatic numbers of some metric spaces," *Russian Mathematical Surveys*, vol. 56, no. 1, pp. 103–139, 2001.

[6] A. M. Raigorodskii, "Coloring distance graphs and graphs of diameters," in *Thirty Essays on Geometric Graph Theory*, pp. 429–460, Springer, 2013.

[7] A. M. Raigorodskii, "Cliques and cycles in distance graphs and graphs of diameters," *Discrete Geometry and Algebraic Combinatorics*, vol. 625, pp. 93–110, 2014.

[8] J. B. Saxe, "Embeddability of weighted graphs in k-space is strongly np-hard," in *Proc. 17th Allerton Conf. Commun. Control Comput*, pp. 480–489, 1979.

[9] B. Horvat, J. Kratochvíl, and T. Pisanski, "On the computational complexity of degenerate unit distance representations of graphs," in *Combinatorial algorithms*, pp. 274–285, Springer, 2011.

[10] L. Lovász, "Self-dual polytopes and the chromatic number of distance graphs on the sphere," *Acta Scientiarum Mathematicarum*, vol. 45, no. 1-4, pp. 317–323, 1983.

[11] A. Raigorodskii, "On the chromatic numbers of spheres in euclidean spaces," in *Doklady Mathematics*, vol. 81, pp. 379–382, Springer, 2010.

[12] A. Raigorodskii, "On the chromatic numbers of spheres in $\mathbb{R}^n$," *Combinatorica*, vol. 32, no. 1, pp. 111–123, 2012.

[13] R. M. Karp, *Reducibility among combinatorial problems*. Springer, 1972.

[14] I. Niven, "Irrational numbers, carus math," *Monographs, John Wiley and Sons Inc*, 1956.